

Gödel type solution in teleparallel gravity

Yu.N. Obukhov*

Institute for Theoretical Physics, University of Cologne, 50923 Köln, Germany

T. Vargas

Instituto de Física Teórica, Universidade Estadual Paulista, Rua Pamplona 145,

01405-900 São Paulo SP, Brazil

Abstract

The stationary cosmological model *without* closed timelike curves of Gödel type is obtained for the ideal dust matter source within the framework of the teleparallel gravity. For a specific choice of the teleparallel gravity parameters, this solution reproduces the causality violating stationary Gödel solution in general relativity, in accordance with the teleparallel equivalent of general relativity. The relation between the axial-vector torsion and the cosmic vorticity is clarified.

Typeset using REVTeX

*On leave from: Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

I. INTRODUCTION

Cosmological models with rotation attract considerable attention since the early papers of Gödel [1]. However, there are two fundamental difficulties of the original Gödel rotating model which made its physical significance problematic: the absence of expansion and the presence of the closed timelike curves. The latter is equivalent to the possibility of violation of causality in such a spacetime. Much of the research on this subject [2–16] was aimed at removing both drawbacks and thereby in constructing a realistic cosmology with rotation and expansion. An overview of the corresponding developments and a comprehensive list of the relevant references is given in [17].

An alternative approach to gravitation is the so called teleparallel gravity (see [18–20] where the basic references on the relevant literature can be found) which corresponds to a gauge theory for the translation group based on the Weitzenböck geometry [21]. In this theory, gravitation is attributed to torsion [22], which plays the role of a force [23], whereas the curvature tensor vanishes identically. Such a spacetime (Weitzenböck manifold) is characterized by a nontrivial tetrad field as the fundamental structure, and this field gives rise to the metric as a by-product. The translational gauge potentials appear as a nontrivial part of the tetrad field, and they induces on spacetime a teleparallel structure which is directly related to the presence of the gravitational field.

In this work we address the above mentioned problems of a rotating world in the framework of the teleparallel gravity theory. The general teleparallelism model is characterized by the three coupling constants, and as a result the structure of the corresponding field equations is less rigid than that of the general relativity theory. Using this fact, one can expect that there exist classes of teleparallel models which are free of one or both difficulties.

More specifically, in the current paper we demonstrate that the completely causal cosmological Gödel type solutions do exist in the appropriate teleparallel gravity models.

The structure of the paper is as follows. In Section II, we review the main features of general teleparallel gravity. In Section III, we obtain the teleparallel Gödel type solution.

Conclusions are presented in Section IV.

II. TELEPARALLEL GRAVITY

In the teleparallel gravity, the spacetime is represented by the Weitzenböck manifold W_4 of distant parallelism. This gravitational theory naturally arises within the gauge approach based on the group of the spacetime translations. In simplified terms (see [24–27] for the advanced discussion of gauge theory of gravity in the framework of nonlinear connections), this approach introduces at each point of the spacetime manifold a tetrad (coframe) field $h^a{}_\mu$. Its inverse (frame) $h_a{}^\nu$ satisfies the condition $h^a{}_\mu h_a{}^\nu = \delta_\mu^\nu$. The tetrad field induces a teleparallel structure on the spacetime which is directly related to the gravitational field, and the Riemannian metric arises as

$$g_{\mu\nu} = \eta_{ab} h^a{}_\mu h^b{}_\nu, \quad (2.1)$$

with the Minkowski matrix $\eta_{ab} = \text{diag}(+1, -1, -1, -1)$. A parallel transport of a tetrad $h^a{}_\mu$ between two neighbouring points is encoded in the covariant derivative

$$\nabla_\nu h^a{}_\mu = \partial_\nu h^a{}_\mu - \Gamma^\alpha{}_{\mu\nu} h^a{}_\alpha, \quad (2.2)$$

where $\Gamma^\alpha{}_{\mu\nu}$ is the Weitzenböck connection. Imposing the geometric condition that the tetrads are parallelly transported in the Weitzenböck space-time, we obtain

$$\nabla_\nu h^a{}_\mu = \partial_\nu h^a{}_\mu - \Gamma^\alpha{}_{\mu\nu} h^a{}_\alpha \equiv 0.$$

This is the condition of the absolute parallelism, or teleparallelism. It is equivalent to

$$\Gamma^\alpha{}_{\mu\nu} = h_a{}^\alpha \partial_\nu h^a{}_\mu \quad (2.3)$$

which gives the explicit form of the Weitzenböck connection in terms of the tetrad. The antisymmetric part of connection,

$$T^\rho{}_{\mu\nu} = \Gamma^\rho{}_{\nu\mu} - \Gamma^\rho{}_{\mu\nu}, \quad (2.4)$$

is the torsion tensor of the Weitzenböck connection. The curvature of the Weitzenböck connection vanishes identically as a consequence of absolute parallelism.

It has been shown in [22] that the most general action in the teleparallel gravity coupled to the matter has the form

$$S = \int d^4x h \left[\frac{1}{4\kappa} (L + \Lambda) + L^{\text{mat}} \right], \quad (2.5)$$

with $h = \det h^a{}_\mu$, and

$$L = c_1 T^{\lambda\tau\nu} T_{\lambda\tau\nu} + c_2 T^{\lambda\tau\nu} T_{\nu\tau\lambda} + c_3 T^{\sigma\lambda}{}_\sigma T^\sigma{}_{\lambda\sigma}. \quad (2.6)$$

Here c_1 , c_2 , and c_3 are the three dimensionless coupling constants, Λ is a cosmological term, and κ is the Einstein gravitational constant. For the specific choice of the parameters,

$$c_1 = \frac{1}{4}, \quad c_2 = \frac{1}{2}, \quad c_3 = -1, \quad (2.7)$$

the teleparallel gravity reduces to the so called teleparallel equivalent of the general relativity.

The Lagrangian (2.6) may be rewritten as

$$L = S^{\lambda\tau\nu} T_{\lambda\tau\nu}, \quad (2.8)$$

where

$$S^{\lambda\tau\nu} = c_1 T^{\lambda\tau\nu} + \frac{c_2}{2} (T^{\tau\lambda\nu} - T^{\nu\lambda\tau}) + \frac{c_3}{2} (g^{\lambda\nu} T^{\sigma\tau}{}_\sigma - g^{\tau\lambda} T^{\sigma\nu}{}_\sigma). \quad (2.9)$$

In the present paper, we use the traditional tensor formalism. Those readers, who prefer a more advanced exterior calculus approach, should address [28] for a general discussion of teleparallel gravity in the metric-affine approach. Here, we only mention that the gravity field Lagrangian can be rewritten in terms of the quadratic contractions of the three irreducible torsion pieces with the corresponding coupling constants a_1, a_2, a_3 . Then one can verify that the original coupling constants c_1, c_2, c_3 are related to the new parameters as follows: $c_1 = (2a_1 + a_3)/3$, $c_2 = 2(a_2 - a_1)/3$, and $c_3 = 2(a_3 - a_1)/3$, see [28].

By performing variation of the action (2.5) with respect to $h^a{}_\mu$, we get the teleparallel gravitational field equation

$$\frac{1}{h} \partial_\sigma (h S_\lambda^{\rho\sigma}) + S_\nu^{\mu\rho} \Gamma^\nu_{\mu\lambda} - \frac{1}{4} \delta_\lambda^\rho L = \kappa T_\lambda^\rho + \Lambda \delta_\lambda^\rho, \quad (2.10)$$

where $T_\lambda^\rho = h^a_\lambda \delta(h L^{\text{mat}})/(h \delta h^a_\rho)$ is the energy-momentum tensor of matter. In the rest of the paper we confine our attention to the case of the ideal fluid with

$$T_\lambda^\rho = (\varepsilon + p) u_\lambda u^\rho - p \delta_\lambda^\rho. \quad (2.11)$$

Here ε is the energy density, p is pressure, and u^μ is the fluid 4-velocity (normalized by $u^\mu u_\mu = 1$).

III. THE TELEPARALLEL GÖDEL TYPE SOLUTION

The line element of the Gödel type cosmological model is given by the interval

$$ds^2 = dt^2 - 2\sqrt{\sigma} a(t) e^{mx} dt dy - a(t)^2 (dx^2 + k e^{2mx} dy^2 + dz^2), \quad (3.1)$$

where m, σ, k are the constant parameters and $a(t)$ is the time-dependent cosmological scale factor. A general analysis [5,17] of the kinematic properties of such a spacetime shows that the closed timelike curves are absent in this manifold when $k > 0$ ($\sigma > 0$ by definition).

Using the relation (2.1) we obtain the tetrad:

$$\tilde{h}^a_\mu = \begin{pmatrix} 1 & 0 & -\sqrt{\sigma} a e^{mx} & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & a\sqrt{\sigma+k} e^{mx} & 0 \\ 0 & 0 & 0 & a \end{pmatrix}, \quad (3.2)$$

and its inverse is

$$\tilde{h}_a^\mu = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & a^{-1} & 0 & 0 \\ \sqrt{\frac{\sigma}{k+\sigma}} & 0 & \frac{e^{-mx}}{a\sqrt{\sigma+k}} & 0 \\ 0 & 0 & 0 & a^{-1} \end{pmatrix}. \quad (3.3)$$

Using the Lorentz transformation for the tetrad $\tilde{h}^a_\mu$

$$h^a_\mu = \Lambda^a_b \tilde{h}^b_\mu, \quad (3.4)$$

we obtain the Weitzenböck connection

$$\Gamma_{\mu\nu}^{\alpha} = \tilde{\Gamma}_{\mu\nu}^{\alpha} + \tilde{h}_b^{\alpha} (\Lambda^{-1} \partial_{\nu} \Lambda)^b_c \tilde{h}^c_{\mu}. \quad (3.5)$$

We assume that the Lorentz matrix depends on the cosmological time only. Hence (denoting the time derivative by the dot) $\partial_{\nu} \Lambda^a_c = \delta_{\nu}^t \dot{\Lambda}^a_c$ and consequently

$$(\Lambda^{-1} \partial_{\nu} \Lambda)^b_c = \delta_{\nu}^t L^b_c, \quad (3.6)$$

with an antisymmetric matrix ($L_{bc} = -L_{cb}$) of the form

$$L^b_c = \begin{pmatrix} 0 & v_1 & v_2 & v_3 \\ v_1 & 0 & -s_3 & s_2 \\ v_2 & s_3 & 0 & -s_1 \\ v_3 & -s_2 & s_1 & 0 \end{pmatrix}. \quad (3.7)$$

From Eqs. (3.2), (3.3) and (3.7), we can now construct the Weitzenböck connection. Its nonvanishing components read:

$$\begin{aligned} \Gamma_{tt}^t &= v_2 \sqrt{\frac{\sigma}{\sigma+k}}, & \Gamma_{xt}^t &= a \left(v_1 + s_3 \sqrt{\frac{\sigma}{\sigma+k}} \right), \\ \Gamma_{yt}^t &= v_2 \frac{ae^{mx}k}{\sqrt{\sigma+k}}, & \Gamma_{zt}^t &= a \left(v_3 - s_1 \sqrt{\frac{\sigma}{\sigma+k}} \right), \\ \Gamma_{tt}^x &= \frac{v_1}{a}, & \Gamma_{xt}^x &= \Gamma_{zt}^z = \frac{\dot{a}}{a}, & \Gamma_{yt}^x &= -e^{mx}(s_3 \sqrt{\sigma+k} + v_1 \sqrt{\sigma}), \\ \Gamma_{zt}^x &= -\Gamma_{xt}^z = s_2, & \Gamma_{tt}^y &= v_2 \frac{e^{-mx}}{a\sqrt{\sigma+k}}, \\ \Gamma_{xt}^y &= s_3 \frac{e^{-mx}}{\sqrt{\sigma+k}}, & \Gamma_{yt}^y &= \frac{\dot{a}}{a} - v_2 \sqrt{\frac{\sigma}{\sigma+k}}, & \Gamma_{zt}^y &= -s_1 \frac{e^{-mx}}{\sqrt{\sigma+k}}, \\ \Gamma_{yx}^y &= m, & \Gamma_{tt}^z &= \frac{v_3}{a}, & \Gamma_{yt}^z &= e^{mx}(s_1 \sqrt{\sigma+k} - v_3 \sqrt{\sigma}). \end{aligned} \quad (3.8)$$

Here, as before, the dot denotes a derivative with respect to the time t .

Assuming that the matter is co-moving with the space, i.e., taking the velocity $u^{\alpha} = \delta_t^{\alpha}$, we find the components of the energy-momentum tensor (2.11):

$$T_t^t = \varepsilon, \quad T_x^x = T_y^y = T_z^z = -p, \quad T_y^t = -\sqrt{\sigma} a e^{mx} (\varepsilon + p). \quad (3.9)$$

A. Solving the field equations

In this paper we will study the case of the stationary world with the constant scale factor $a = \text{const}$. In this way we will be able to perform a direct comparison of the solutions obtained with the Gödel solution in the general relativity theory.

The value of the parameter $2c_1 - c_2$ is essential. When this is equal zero, i.e., $2c_1 = c_2$ [note that this case includes the teleparallel equivalent of general relativity (2.7)], then the difference of the $(_y^y)$ and $(_z^z)$ components of the field equation (2.10) yields:

$$\frac{(2k + \sigma) m c_2}{2a^2(k + \sigma)} = 0. \quad (3.10)$$

In other words, we have three possibilities: either $m = 0$, then the cosmic vorticity vanishes, or the coupling constant $c_2 = 0$, but then also $c_1 = 0$ and the resulting Lagrangian is unphysical. Finally, when both m and c_2 are nontrivial, we should have $k = -\sigma/2$. This is exactly the value of the parameter k of the original Gödel model which contains the closed timelike curves. Accordingly, we will not consider this case as it basically leads to the qualitatively same model.

Thus we instead assume that $2c_1 - c_2 \neq 0$. Then the $(_z^t)$ -th component of the field equation yields $s_2 = 0$, whereas $(_x^z)$ -th component yields

$$s_1 = v_3 \sqrt{\frac{\sigma}{\sigma + k}}. \quad (3.11)$$

Furthermore, subtracting $(_y^y)$ from $(_z^z)$, we find

$$s_3 = -v_1 \sqrt{\frac{\sigma}{\sigma + k}} + \frac{m}{a} \frac{2kc_1 + (k + \sigma)c_2}{(2c_1 - c_2)\sqrt{\sigma}\sqrt{\sigma + k}}. \quad (3.12)$$

Now, the component $(_t^t)$ minus $(_x^x)$ plus the product of $(e^{-mx}/a\sqrt{\sigma})$ times $(_y^t)$ yields the equation

$$\frac{k\sigma(2c_1 + c_2 + c_3)[k(v_1^2 + v_3^2) + (k + \sigma)v_2^2] + v_1 \frac{m}{a} Q_1 + 2 \frac{m^2}{a^2} P_1}{2\sigma(k + \sigma)^2} = 0. \quad (3.13)$$

Here we have denoted the following combinations of the constants:

$$Q_1 = \frac{k\sigma(2c_1 + c_2 + c_3)[2(k - \sigma)c_1 + 3(k + \sigma)c_2]}{2c_1 - c_2}, \quad (3.14)$$

$$P_1 = [\sigma c_1 - (k + \sigma)c_2][4(2k^2 + 2k\sigma + \sigma^2)c_1^2 - 2\sigma(2k + \sigma)c_1c_2 - 2k\sigma c_1c_3 - 2(k + \sigma)^2c_2^2 - \sigma(k + \sigma)c_2c_3]/(2c_1 - c_2)^2. \quad (3.15)$$

The (t^x) -th and the (t^y) -th components of the field equation read, respectively:

$$v_2 m \frac{(2c_1 + c_2)[2(k + \sigma)c_1 - (3k + 2\sigma)c_2] - [2kc_1 + (k + \sigma)c_2]c_3}{2\sqrt{\sigma}\sqrt{k + \sigma}(2c_1 - c_2)a^2} = 0, \quad (3.16)$$

$$\frac{-\sigma(2c_1 + c_2 + c_3)[k(v_1^2 + v_3^2) + (k + \sigma)v_2^2] - v_1 \frac{m}{a}Q_2 + 2\frac{m^2}{a^2}P_2}{2\sqrt{\sigma}e^{mx}a(k + \sigma)^2} = 0. \quad (3.17)$$

Here we have introduced further abbreviations:

$$Q_2 = \left\{ (2c_1 + c_2) [2k(k + 3\sigma)c_1 - (3k - \sigma)(k + \sigma)c_2 - (k + \sigma)^2c_3] + [4kc_1 + 3(k + \sigma)c_2]c_3\sigma \right\} / (2c_1 - c_2), \quad (3.18)$$

$$P_2 = \frac{(2c_1 + c_2 + c_3)[2\sigma c_1 + (k + \sigma)c_2][(k + \sigma)c_2 - \sigma c_1]}{(2c_1 - c_2)^2}. \quad (3.19)$$

Let us analyse the two equations (3.13) and (3.17) which can be considered as the algebraic system for v_1 and v_3 . Then discarding the denominators, from the difference and the sum of the numerators in (3.13) and (3.17) we find

$$v_1 = -\frac{2m}{a} \frac{P_1 + kP_2}{Q_1 - kQ_2} = \frac{2m[\sigma c_1 - (k + \sigma)c_2]}{a k(2c_1 - c_2)}, \quad (3.20)$$

$$\begin{aligned} \sigma(2c_1 + c_2 + c_3)[k(v_1^2 + v_3^2) + (k + \sigma)v_2^2] &= \frac{2m^2}{a^2} \frac{P_1Q_2 + P_2Q_1}{Q_1 - kQ_2} \\ &= -\frac{2m^2P_0[\sigma c_1 - (k + \sigma)c_2]}{a^2 k(2c_1 - c_2)^2}. \end{aligned} \quad (3.21)$$

Here we denoted

$$\begin{aligned} P_0 &= 4k(2k + 3\sigma)c_1^2 + 2\sigma(2k + \sigma)c_1c_2 - \sigma^2c_1c_3 \\ &\quad - (k + \sigma)(2k - \sigma)c_2^2 + 2\sigma(k + \sigma)c_2c_3. \end{aligned} \quad (3.22)$$

Note that although the expressions (3.14), (3.15), and (3.18), (3.19) are quite nontrivial, the final result (3.20)-(3.21) is rather simple. Substituting (3.20) into (3.21), we obtain explicitly

$$v_3^2 = \frac{k + \sigma}{k} \left[-v_2^2 + \frac{2m^2(2k + \sigma) [(k + \sigma)c_2 - \sigma c_1](2c_1 + c_2)}{a^2 k \sigma (2c_1 + c_2 + c_3)(2c_1 - c_2)} \right]. \quad (3.23)$$

Obviously, we can put $v_2 = 0$ from the equation (3.16). Otherwise (3.16) determines the parameter k in terms of σ and c_1, c_2, c_3 .

Finally, taking into account (3.11), (3.12), (3.20), and (3.23), we can write down the diagonal components of the field equation (2.10):

$$-\frac{3m^2k(2c_1 - c_2)}{2\sigma a^2} = \kappa \varepsilon + \Lambda, \quad (3.24)$$

$$-\frac{m^2k(2c_1 - c_2)}{2\sigma a^2} = -\kappa p + \Lambda. \quad (3.25)$$

In order to compare with the old Gödel model, we now specialise to the case of the dust cosmological matter, $p = 0$. Then equation (3.25) determines the parameter k/σ in terms of the coupling and geometrical constants, whereas (3.24) yields the energy density of the stationarily rotating world:

$$\varepsilon = \frac{m^2k(c_2 - 2c_1)}{\kappa \sigma a^2}. \quad (3.26)$$

Evidently, for the positivity of energy, the teleparallel coupling constants should satisfy the inequality $c_2 > 2c_1$. The important point is that the ratio k/σ must also be then positive. Hence the resulting rotating universe does not contain *closed* timelike curves, and consequently the causality is not violated.

B. Finding the Lorentz matrix factor

In order to complete the description of the exact solution obtained, it remains to find the Lorentz matrix Λ^a_b which enters the tetrad (3.4). It should be noted that from the point of view of the resulting Riemannian geometry, this Lorentz factor is irrelevant since the spacetime interval (3.1) is invariant under the local Lorentz transformations of the tetrad. However, since we deal with the Weitzenböck geometry, the Lorentz factor is the essential element of the definition of the tetrad (3.4) and, accordingly, of the connection

(3.8). Assuming that the elements of the unknown matrix Λ^a_b depend only on cosmological time t , we can rewrite the differential equation (3.6) as

$$\dot{\Lambda}^a_b = \Lambda^a_c L^c_b. \quad (3.27)$$

Fixing the index a , we thus find that the four elements of each row Λ^0_a , or Λ^1_a , or Λ^2_a , or Λ^3_a satisfy the system of the linear differential equations (3.27) with the constant coefficients

$$L^a_b = \begin{pmatrix} 0 & v_1 & 0 & v_3 \\ v_1 & 0 & -s_3 & 0 \\ 0 & s_3 & 0 & -s_1 \\ v_3 & 0 & s_1 & 0 \end{pmatrix}. \quad (3.28)$$

The elements of this matrix were determined above in the equations (3.11), (3.12), (3.20) and (3.23) and they depend on the coupling constants of the model and on the geometric parameters of the ansatz. However, the explicit form of these relations is not needed for our purposes. The solution of such a system is straightforward: We assume the standard ansatz of the $\Lambda^b_a = e^{\lambda t} \alpha^b_a$. Then the constant coefficients α^b_a are nontrivial, provided the characteristic equation is satisfied:

$$\begin{vmatrix} -\lambda & v_1 & 0 & v_3 \\ v_1 & -\lambda & s_3 & 0 \\ 0 & -s_3 & -\lambda & s_1 \\ v_3 & 0 & -s_1 & -\lambda \end{vmatrix} = 0. \quad (3.29)$$

Evaluating the determinant, we find the algebraic equation

$$\lambda^4 + \lambda^2 (s_1^2 + s_3^2 - v_1^2 - v_3^2) - (s_1 v_1 + s_3 v_3)^2 = 0. \quad (3.30)$$

It is easy to see that of the four roots $\lambda_{(A)}$, $A = 1, \dots, 4$, two are real and two are purely imaginary. When λ is a solution of the equation (3.30), the components α^a_0 and α^a_1 represent the independent free parameters, whereas the rest of the components are expressed as

$$\alpha^a_2 = \frac{\lambda^2 - v_1^2 - v_3^2}{v_1 s_3 - v_3 s_1} \alpha^a_0, \quad \alpha^a_3 = \frac{\lambda^2 + s_3^2 - v_1^2}{s_1 s_3 + v_1 v_3} \alpha^a_1. \quad (3.31)$$

The general solution of the system (3.27) has the form

$$\Lambda^a_b = \sum_{i=A}^4 C_{(A)} \alpha^a_b e^{\lambda_{(A)} t}, \quad (3.32)$$

where the $C_{(A)}$ are arbitrary constants. It is thus interesting to note that whereas the geometry of the solution is stationary, with the Riemannian curvature and the Weitzenböck torsion tensors being independent of the cosmological time, the teleparallel tetrad has a nontrivial time dependence via the Lorentz matrix (3.32), in general.

IV. DISCUSSION AND CONCLUSION

In this paper we have obtained the teleparallel version of the stationary rotating cosmological solution of Gödel type within the framework of the 3-parameter model with the general teleparallel Lagrangian (2.6). In particular, we have demonstrated that even for the ideal dust matter source there exists a solution *without* closed timelike curves and thus the causality is not violated in such a spacetime.

For a specific choice of the teleparallel gravity parameters, this solution reproduces the causality violating stationary Gödel solution in general relativity, in accordance with the teleparallel equivalent of general relativity.

Previously it was conjectured that the axial torsion is directly related to the spacetime rotation, showing that such an assumption is valid for the case of the Kerr geometry [29]. It is interesting to check whether this is also true for the rotating cosmology under consideration.

Using the decomposition of the torsion into three irreducible parts under the group of global Lorentz transformations [22], we can straightforwardly calculate the axial-vector part

$$A^\mu = \frac{1}{6} \eta^{\mu\nu\rho\sigma} T_{\nu\rho\sigma}. \quad (4.1)$$

Here $\eta_{\mu\nu\rho\sigma}$ is the completely antisymmetric Levi-Civita tensor. Explicitly, we find from (3.8):

$$A^t = 0, \quad A^x = -\frac{2}{3a} \left(s_1 - v_3 \sqrt{\frac{\sigma}{k+\sigma}} \right), \quad A^y = -\frac{2}{3a} \frac{e^{-mx}}{\sqrt{k+\sigma}} s_2, \quad (4.2)$$

$$A^z = -\frac{2}{3a} \left(s_3 + v_1 \sqrt{\frac{\sigma}{k+\sigma}} \right) - \frac{m}{3a^2} \sqrt{\frac{\sigma}{\sigma+k}}. \quad (4.3)$$

For the exact solution obtained, using the equations (3.11) and (3.12), we prove that the first three components vanish $A^t = A^x = A^y = 0$, whereas

$$A^z = -\frac{m(2c_1 + c_2)(\sigma + 2k)}{3a^2(2c_1 - c_2)\sqrt{\sigma}\sqrt{k + \sigma}}. \quad (4.4)$$

As we can see, the axial vector torsion is, indeed, related to the spacetime vorticity. In particular, the axial torsion is apparently directed along the axis of the cosmic rotation and its value is proportional to the magnitude of the vorticity. However, this relation should not be overestimated. Let us recall that $k = -\sigma/2$ in the original Gödel model. Then we find from (4.4) that the axial torsion vanishes completely, $A^\mu = 0$, for this case. The existence of the rotating cosmological models with the vanishing axial torsion shows that one cannot reduce the description of the cosmic vorticity to the properties of the irreducible parts of the Weitzenböck torsion tensor.

The earlier studies of the teleparallel gravity have revealed the one-parameter family of models which are compatible with the observational data [22,19,31,32]. Another characteristic feature of that family is the existence of black hole solutions which seem to be absent in the generic case [28]. On the other hand, exactly this family appears to be plagued with the problem of ghost and negative-energy modes as well as of the “hidden” gauge transformations [19,30–32], see also a recent study in [33]. In this sense, it is interesting that the nontrivial spatial axial-vector torsion (4.1) of our solution prevents the occurrence of the special “hidden” gauge transformation [31] of the torsion. At the same time, it is worthwhile to note that the one-parameter family admits only the causality violating cosmological solutions with rotation, whereas the timelike closed curves are absent in the generic case.

Acknowledgments. For YNO this work was supported by the Deutsche Forschungsgemeinschaft (Bonn) with the grant HE 528/20-1. TV would like to thank FAPESP grant 02/04199-1 for financial support.

REFERENCES

- [1] K. Gödel, Rev. Mod. Phys. 21 (1949) 447.
- [2] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Space–Time* (Cambridge University Press, Cambridge, 1973).
- [3] M.J. Rebouças, Phys. Lett. A 70 (1979) 161.
- [4] J. Pfarr, Gen. Relat. Grav. 13 (1981) 1073.
- [5] S. Maitra, J. Math. Phys. 7 (1966) 1025.
- [6] I. Ozsváth and E. Schücking, Ann. Phys. 55 (1969) 166.
- [7] P. Birch, Nature 298, 1982, 451.
- [8] B. Nodland and J.P. Ralston, Phys. Rev. Lett. 78 (1997) 3043.
- [9] L.X. Li, Gen. Relat. Grav. 30 (1998) 497.
- [10] A.J. Aciolly, Progr. Theor. Phys. 79 (1988) 1269.
- [11] J. Hwang and H. Noh, Phys. Rev. D 57 (1998) 2617.
- [12] J.G. Williams, Gen. Relat. Grav. 26 (1994) 499.
- [13] M. Bedran, V. Vaidya, and M. Som, Nuovo Cim. B 87 (1985) 101.
- [14] L.L. Smalley, Phys. Lett. A 113 (1986) 463.
- [15] Yu.N. Obukhov and V.A. Korotky, Class. Quantum Grav. 4 (1987) 1633.
- [16] J.E. Aman, J.B. Fonseca-Neto, M.A.H. MacCallum, and M.J. Rebouças, Class. Quantum Grav. 15 (1998) 1089.
- [17] Yu.N. Obukhov, *On physical foundations and observational effects of cosmic rotation*, in: “Colloquium on Cosmic Rotation” (Berlin, Feb. 1998), Eds. M. Scherfner, T. Chrobok, and M. Shefaat (Wissenschaft und Technik Verlag: Berlin, 2000) 23-96; E-print archive:

astro-ph/0008106.

- [18] F. Gronwald and F.W. Hehl, *On the gauge aspects of gravity*, in: *International School of Cosmology and Gravitation: 14th Course: “Quantum Gravity”*, May 1995, Erice, Italy. Proceedings. P.G. Bergmann, V. de Sabbata, and H.J. Treder (eds.) (World Scientific: Singapore, 1996) pp. 148–198. E-print archive: gr-qc/9602013.
- [19] F. Mueller-Hoissen and J. Nitsch, Phys. Rev. D 28 (1983) 718.
- [20] V.C. de Andrade, L.C.T. Guellin, and J.G. Pereira, *Teleparallel Gravity: An Overview*, in: Proceedings of the IX Marcel Grossmann’s Meeting, Rome, Italy, July, 2000; E-print archive: gr-qc/0011087.
- [21] R. Weitzenböck, *Invariantentheorie* (Gronningen: Noordhoff, 1923).
- [22] K. Hayashi and T. Shirafuji, Phys. Rev. D 19 (1978) 3524.
- [23] V.C. de Andrade and J. G. Pereira, Phys. Rev. D 56 (1997) 4689.
- [24] V.N. Ponomarev, A.O. Barvinski, and Yu.N. Obukhov, *Geometrodynamical methods and the gauge approach to the theory of gravitational interactions* (Energoatomizdat: Moscow, 1985), in Russian.
- [25] R. Percacci, *Geometry of nonlinear field theories* (World Scientific: Singapore, 1986).
- [26] F. Gronwald, Intern. J. Mod. Phys. D6 (1997) 263.
- [27] R. Tresguerres, Phys. Rev. D66 (2002) 064025.
- [28] Yu.N. Obukhov and J.G. Pereira, Phys. Rev. D67 (2003) 044016.
- [29] J.G. Pereira, T. Vargas, and C.M. Zhang, Class. Quantum Grav. 18 (2001) 833.
- [30] W. Kopczyński, Phys. A15 (1982) 493.
- [31] J.M. Nester, Class. Quantum Grav. 5 (1988) 1003.

- [32] R.D. Hecht, J. Lemke, and R.P. Wallner, Phys. Rev. D44 (1991) 2442.
- [33] M. Blagojevic, M. Vasilic, Class. Quantum Grav. 17 (2000) 3785.